






ACL Reconstruction with Bone-Patellar Tendon-Bone (BTB) Autograft and All Inside Meniscal Repair Rehabilitation Protocol

	Weight-bearing	Brace	Range of Motion	Exercises
Phase 1 0-6 weeks	Full in brace ^a	0-2 week: Locked in full extension for ambulation and sleeping Off for exercises and hygiene only 2-6 weeks: Unlocked 0-90° for ambulation, remove for sleeping	Passive 0-90° at one week postop visit, then progress to full	Calf pumps, patellar mobilization, quadriceps tightening sets and SLR in brace . No weight-bearing motion. Add side-lying hip and core, advance quad set and stretching ^a In brace: Addition of heel raises, total gym (closed chain), gait normalization, eccentric quads, eccentric hamstrings. Advance core work with hip, gluteal and core strengthening No weight-bearing with flexion >90°
Phase 2 6-12 weeks	Full, progressing to normal gait pattern	Discontinue at 6 weeks if quadriceps control is adequate (no extension lag)	Gain full and pain-free	Begin toe raises, closed chain quads, balance exercises, hamstring curls, stationary bike, step-ups, front and side planks, hip/glute/core
Phase 3 12-16 weeks	Full with normalized gait pattern	None	Full	Advance closed chain strengthening Progress proprioception activities Begin stair climber, elliptical at 8 weeks and running straight at 12 weeks when 8" step down is satisfactory
Phase 4 16 weeks – 6 months	Full	None	Full	16 weeks: Begin jumping 20 weeks: Advance running to sprinting, backward running, cutting, pivoting, changing direction, initiate plyometric program and sport-specific drills 22 weeks: Advance as tolerated FSA completed at 22 weeks ^b
Phase 5 6-12 months	Full	None	Full	Gradual return to sports participation after completion of functional sports assessment (FSA) Maintenance program based on FSA

SLR – Straight-leg raise

^aModified with concomitantly performed meniscus repair/transplantation or articular cartilage procedure

^bCompletion of FSA (Functional Sports Assessment) is not mandatory, but is recommended at approximately 22 wks post-op for competitive athletes returning to play after rehabilitation

Jason R. Ferrel, MD | 614-545-7958 |      @JasonFerrel_MD